

Inverse problems for the number of maximal independent sets

Alex Dainiak

November 21, 2011

Abstract

We study the following inverse graph-theoretic problem: how many vertices should a graph have given that it has a specified value of some parameter. We obtain asymptotic for the minimal number of vertices of the graph with the given number n of maximal independent sets for a class of natural numbers that can be represented as concatenation of periodic binary words.

Problems of estimating various graph invariants play the central role in quantitative graph theory. Among the most studied invariants are connectivity, chromatic number, girth, independence number, maximal clique size, number of independent sets etc. As well as forward problems, inverse problems also are of interest. They generally can be stated as follows: find a graph (or prove its existence) that have the desired value of some parameter. The classical problem of this kind is finding a graph with the given degree sequence [2, 3]. For a long time it was not known if there was only a finite number of naturals not being the Wiener index of trees. [5, 6]. An analogous question considering the number of independent sets in trees, asked in [4], is not yet solved, whereas some other parameters of trees are better studied (e.g. [1]).

We now state the problems coeverted in this paper in their general form. Let $\mathcal{G}$ be a family of graphs, and let $\phi : \mathcal{G} \rightarrow S$ and $\psi : \mathcal{G} \rightarrow T$ be arbitrary functionals on $\mathcal{G}$. The *existential inverse problem* for the pair $(\mathcal{G}, \phi)$ may be stated as follows: “describe all $s \in S$ for which there exists a graph $G \in \mathcal{G}$ having $\phi(G) = s$ ”.

Let S be the set of all values of ψ for all graphs in $\mathcal{G}$. For $S \subseteq \mathbb{N}$ we call $\mathcal{G}$ to be *strongly ϕ -complete*, if for every $s \in S$ there is $G \in \mathcal{G}$ such that $\phi(G) = s$. If such $G \in \mathcal{G}$ exists for all large enough $s \in S$, then we say that $\mathcal{G}$ is *weakly ϕ -complete*, or just *ϕ -complete*. If $\phi(G) = s$ then we say that s is *realized* by G .

If the existential inverse problem is solved positively, we can consider the *optimizational inverse problem* for the triple $(\mathcal{G}, \phi, \psi)$: “for a given $s \in S$ find $L_{\phi, \psi}^{\mathcal{G}}(s) = \inf\{\psi(G) \mid G \in \mathcal{G}, \phi(G) = s\}$ ”. As the problem of finding L exactly is too hard, it is natural to consider only the asymptotic behavior of $L_{\phi, \psi}^{\mathcal{G}}(s)$ for ϕ -complete families of graphs. If $\mathcal{G}$ is a class of all graphs, we shorten the notation $L_{\phi, \psi}^{\mathcal{G}}(s)$ to $L_{\phi, \psi}(s)$.

Denote by $\iota(G)$ the number of all independent sets (i. s.) of vertices in G , and by $\iota_m(G)$ the number of maximal-by-inclusion i.s. (m. i. s.) in G . Finally, by $\iota_M(G)$ we denote the number of maximum independent sets in G . We write $\nu(G)$ and $\epsilon(G)$ for the number of vertices and edges in G respectively. The families of bipartite graphs and forests are denoted by $\mathcal{B}$ and $\mathcal{F}$ respectively. We write K_r and P_r for complete graphs and paths on r vertices. $K_{r,s}$ denotes complete bipartite graph, r and s being the sizes of its parts. $K'_{r,r}$ stands for the *corona-graph*, which can be constructed by deleting edges of some perfect matching from $K_{r,r}$. The sets of

vertices and edges of graph G will be denoted as V_G and E_G . The edge between vertices u and v will be denoted as uv . A family of all m.i.s. of G is denoted as $\mathcal{I}_m(G)$.

In the notation of the current paper, Linek proved [4] the strong ι -completeness of $\mathcal{B}$. It justifies the consideration of the optimization problem of finding $L_{\iota, \nu}^{\mathcal{B}}(n)$. The trivial lower bound is $L_{\iota, \nu}^{\mathcal{B}}(n) \geq \log_2 n$ (which follows from the inequation $\iota(G) \leq 2^{\nu(G)}$). A graph constructed in [4] to realize a given natural number has the maximal possible sizes of parts: $\lfloor \log_2 n \rfloor$ and $\lfloor \log_2(n - 2^{\lfloor \log_2 n \rfloor + 1}) \rfloor$. For $n = 2^k - 1$ such graph would have $2k - 2$ vertices, which is double the expected optimal size. Some n of the above form can be realized more economically, as the following statement shows.

Theorem 1. *For $k = 2^t$ we have $L_{\iota, \nu}^{\mathcal{F}}(2^k - 1) \lesssim k$.*

Proof. Just note that

$$2^{2^t} - 1 = \prod_{j=0}^{t-1} (2^{2^j} + 1) = \prod_{j=0}^{t-1} \iota(K_{2^j, 1}) = \iota\left(\bigsqcup_{j=0}^{t-1} K_{2^j, 1}\right).$$

At the same time $\nu(\bigsqcup_{j=0}^{t-1} K_{2^j, 1}) = 2^t + t - 1 \lesssim k$. $\square$

The existential inverse problem for $(\mathcal{B}, \iota_m)$ is trivial, as any $n \geq 4$ can be realized as the number of m.i.s. in corona-graph $K'_{n-2, n-2}$. If we consider ψ to be the number of vertices of a graph, we come up with an optimizational inverse problem: “for natural n find minimal $L(n)$ such that there exists a graph on $L(n)$ vertices having n maximal independent sets”. The remaining part of the paper is dedicated to estimating $L(n)$.

Bounds for $L_{\iota_m, \nu}^{\mathcal{B}}(n)$

Lemma 1. *Let G be bipartite with parts L_G, R_G and without isolated vertices. Let $\tilde{G}$ be a bipartite graph, vertex-disjoint with G , and let U_1 and U_2 be some subsets of first and second parts of $\tilde{G}$ respectively. Let G' be a graph obtained by connecting all vertices in U_1 (resp. U_2) to all vertices in L_G (resp. R_G). Then we have*

$$\iota_m(G') = (\iota_m(G) - 2) \cdot \iota_m(\tilde{G} \setminus (U_1 \cup U_2)) + \iota_m(\tilde{G} \setminus U_1) + \iota_m(\tilde{G} \setminus U_2) + \iota_m(\tilde{G} + U_1 + U_2),$$

where $\iota_m(\tilde{G} + U_1 + U_2)$ stands for the number of m.i.s. of $\tilde{G}$ having non-empty intersections with both U_1 and U_2 .

Proof. The statement of the lemma can be checked by direct counting. If an m.i.s. of G' contains no vertices of G , then it must contain at least one vertex from both U_1 and U_2 , the number of such sets being $\iota_m(\tilde{G} + U_1 + U_2)$. If a m.i.s. of G' contains vertices from both parts of G , then it is disjoint with $U_1 \cup U_2$, and its subsets in G and $\tilde{G}$ must themselves be maximal independent sets in G and $\tilde{G}$ respectively. Thus the number of such m.i.s. equals $(\iota_m(G) - 2) \cdot \iota_m(\tilde{G} \setminus (U_1 \cup U_2))$. If a m.i.s. of G contains all vertices of L_G or whole R_G , then its subset in $\tilde{G}$ will form a m.i.s. in $\tilde{G} \setminus U_1$ or $\tilde{G} \setminus U_2$ respectively. $\square$

Let $\tilde{G}$ be bipartite with U_1 and U_2 being some subsets of its parts. Put

$$h'_{\tilde{G}} = \iota_m(\tilde{G} \setminus (U_1 \cup U_2)),$$

$$h''_{\tilde{G}} = (\iota_m(\tilde{G} \setminus U_1) + \iota_m(\tilde{G} \setminus U_2) + \iota_m(\tilde{G} + U_1 + U_2) - 2\iota_m(\tilde{G} \setminus (U_1 \cup U_2))).$$

Lemma 2. *Let Γ be a finite set of bipartite graphs with selected subsets in their parts, such that $\{h'_{\tilde{G}}k + h''_{\tilde{G}} \mid k \in \mathbb{N}, \tilde{G} \in \Gamma\} \supseteq (\mathbb{N} \setminus [1, n_0])$ for some n_0 . Put*

$$\gamma = \max \left\{ (\log_2 h'_{\tilde{G}})^{-1} \nu(\tilde{G}) \mid \tilde{G} \in \Gamma \right\}.$$

Then $L_{\iota_m, \nu}^{\mathcal{B}}(n) \leq \gamma \cdot \log_2 n + O(1)$.

Proof. The lemma is proved by induction on n with a help of lemma 1. Let Γ_0 be an arbitrary finite set of bipartite graphs having $\{\iota_m(G) \mid G \in \Gamma_0\} \supseteq [1, n_0]$. For example, as Γ_0 we can take the set $\{K'_{n-2, n-2} \mid n \in [4, n_0]\} \cup \{K_1, K_{1,1}, P_4\}$. Let ν_0 be the maximal number of vertices of graphs in Γ_0 . It suffices to prove that for any n the following inequality holds:

$$L_{\iota_m, \nu}^{\mathcal{B}}(n) \leq \gamma \cdot \log_2 n + \nu_0, \quad (1)$$

which would imply the statement of the lemma.

The inequality (1) trivially holds for $n \leq n_0$. Consider an arbitrary n' , $n' > n_0$, and assume that (1) holds for all n less than n' . By the conditions of the lemma, there exists some $\tilde{G} \in \Gamma$ and some natural k , such that $n' = h'_{\tilde{G}}k + h''_{\tilde{G}}$. By the induction hypothesis there is some bipartite G having $\iota_m(G) = k$ and $\nu(G) \leq \gamma \cdot \log_2 k + \nu_0$. By lemma 1 we conclude that there is a graph G' with $\iota_m(G') = n'$ and

$$\nu(G') \leq \nu(G) + \nu(\tilde{G}) \leq \nu(\tilde{G}) + \gamma \cdot \log_2 k + \nu_0. \quad (2)$$

By (2) and $k \leq \frac{n'}{h'_{\tilde{G}}}$ we have

$$\begin{aligned} \nu(G') &\leq \nu(\tilde{G}) + \gamma \cdot \log_2 n' - \gamma \cdot \log_2 h'_{\tilde{G}} + \nu_0 = \\ &= \gamma \cdot \log_2 n' + \nu_0 + ((\log_2 h'_{\tilde{G}})^{-1} \nu(\tilde{G}) - \gamma) \cdot \log_2 h'_{\tilde{G}} \leq \\ &\leq \gamma \cdot \log_2 n' + \nu_0. \end{aligned}$$


□

Theorem 2. *For all $n \in \mathbb{N}$ we have*

$$2 \log_2 n \leq L_{\iota_m, \nu}^{\mathcal{B}}(n) \leq 2.88 \log_2 n + O(1). \quad (3)$$

Proof. The lower bound of (3) follows from the observation that a number of m.i.s. in a bipartite graphs cannot exceed the number of subsets of any of this graph's parts.

To obtain the upper bound we apply lemma 2 with Γ being equal to the following set of graphs (subsets U_1, U_2 are marked as bold vertices; pairs of numbers $(h'_{\tilde{G}}, h''_{\tilde{G}})$ are scribed under the graphs):



It can be checked, that such Γ meets the conditions of lemma 2 and for this set the parameter γ would equal $12(\log_2 18)^{-1} < 2.88$. It implies the lower bound in (3). $\square$

Remark. The inequality (3) remains valid without $O(1)$ summand, which can be proven in the same way as in theorem 2. Upper bound in (3) may be directly improved by finding a better set Γ . To find such Γ one can apply an exhaustive computer search (which in fact was used to find Γ that we provide above).

We feel certain that the following is true:

Conjecture. $L_{\iota_m, \nu}^{\mathcal{B}}(n) \sim 2 \log_2 n$ for $n \rightarrow \infty$.

Though we were unable to prove the above conjecture, theorem 3 approves it for some special class of naturals. Next we need to prove some auxillary statements.

Lemma 3. For any bipartite G without isolated vertices there is a bipartite graph without isolated vertices having $(\nu(G) + 4)$ vertices and $(2\iota_m(G) + 1)$ maximal independent sets.

Proof. Apply lemma 1, taking P_4 for $\tilde{G}$, and taking any central vertex of $\tilde{G}$ and empty set for U_1 and U_2 respectively. $\square$

Lemma 4. For any bipartite G without isolated vertices there is a bipartite graph without isolated vertices having $(\nu(G) + 4)$ vertices and $(\iota_m(G) + 2)$ maximal independent sets.

Proof. Apply lemma 1, taking P_4 for $\tilde{G}$, and taking pair of non-adjacent vertices of $\tilde{G}$ and an empty set for U_1 and U_2 respectively. $\square$

Lemma 5. For any bipartite graphs G and $\tilde{G}$ without isolated vertices there is a bipartite graph without isolated vetices with $(\nu(G) + \nu(\tilde{G}) + 4)$ vertices and $(\iota_m(G) + \iota_m(\tilde{G}))$ maximal independent sets.

Proof. Apply lemma 1, with whole parts of $\tilde{G}$ selected as U_1 and U_2 . Thus we obtain G' on $(\nu(G) + \nu(\tilde{G}) + 4)$ vertices with $(\iota_m(G) + \iota_m(\tilde{G}) - 2)$ maximal independent sets. It suffices to apply lemma 4 to G' . $\square$

Lemma 6. *Let G and $\tilde{G}$ be bipartite without isolated vertices, and let $s, t \in \mathbb{N}$. Then there exists bipartite graph without isolated vertices having*

$$2^{st} \cdot \iota_m(G) + \frac{2^{st} - 1}{2^t - 1} \cdot \iota_m(\tilde{G})$$

maximal independent sets and no more than $\nu(G) + \nu(\tilde{G}) + 2s(t + 1) + 3$ vertices.

Proof. For $s = 1$ the statement follows from lemma 5 (before applying the lemma add matching on $2t$ vertices to G). So for the rest of the proof we assume that $s \geq 2$. We also assume that $V_G \cap V_{\tilde{G}} = \emptyset$. Parts of G and $\tilde{G}$ will be denoted as L_G, R_G and $L_{\tilde{G}}, R_{\tilde{G}}$ respectively. We shall consider a graph G' which is constructed as follows:

$$\begin{aligned} V_{G'} &= V_G \cup V_{\tilde{G}} \cup \{w\} \cup \{\tilde{u}_i \mid 1 \leq i \leq t\} \cup \{\tilde{v}_i \mid 1 \leq i \leq t\} \cup \\ &\quad \cup \{u_{i,j} \mid 1 \leq i \leq s-1, 1 \leq j \leq t+1\} \cup \{v_{i,j} \mid 1 \leq i \leq s-1, 1 \leq j \leq t+1\}, \\ E_{G'} &= E_G \cup E_{\tilde{G}} \cup \{\tilde{u}_i \tilde{v}_i \mid 1 \leq i \leq t\} \cup \{u_{i,j} v_{i,j} \mid 1 \leq i \leq s-1, 1 \leq j \leq t\} \cup \\ &\quad \cup \{uv \mid u \in L_G, v \in R_{\tilde{G}}\} \cup \{uv \mid u \in R_G, v \in L_{\tilde{G}}\} \cup \{wv \mid v \in R_G \cup R_{\tilde{G}}\} \cup \\ &\quad \cup \{\tilde{u}_i v \mid 1 \leq i \leq t, v \in R_{\tilde{G}}\} \cup \{u \tilde{v}_i \mid 1 \leq i \leq t, u \in L_{\tilde{G}}\} \cup \\ &\quad \cup \{u_{i,t+1} v \mid 1 \leq i \leq s-1, v \in R_G\} \cup \{u v_{i,t+1} \mid 1 \leq i \leq s-1, u \in L_G\} \cup \\ &\quad \cup \{u_{i,j} v_{k,t+1} \mid 1 \leq i \leq k \leq s-1, 1 \leq j \leq t+1\}. \end{aligned}$$

It can be checked that G' is bipartite with one of its parts being

$$L_{G'} = L_G \cup L_{\tilde{G}} \cup \{w\} \cup \{\tilde{u}_i \mid 1 \leq i \leq t\} \cup \{u_{i,j} \mid 1 \leq i \leq s-1, 1 \leq j \leq t+1\}.$$

We now count all maximal independent sets in G' . These can be of the following seven types:

1. Let $\mathcal{I}_1 = \{I \in \mathcal{I}_m(G') \mid I \cap L_G \neq \emptyset, I \cap R_G \neq \emptyset\}$. Note that for every set $I \in \mathcal{I}_1$ the subset $I \cap V_G$ is m.i.s. in G , and that the intersection of $V_{G'} \setminus V_G$ and I can only contain the following vertices: $\tilde{u}_i, \tilde{v}_i, u_{i,j}$ and $v_{i,j}$ for $j \neq t+1$. The subgraph generated by these vertices is a matching, which implies

$$|\mathcal{I}_1| = (\iota_m(G) - 2) \cdot 2^{st}. \quad (4)$$

2. Let $\mathcal{I}_2 = \{I \in \mathcal{I}_m(G') \mid I \cap L_G \neq \emptyset, I \cap R_G = \emptyset\}$. It can be checked that every $I \in \mathcal{I}_2$ must contain *all* vertices from L_G , and also vertices w and $u_{i,t+1}$ for all i . Moreover I is disjoint with $R_{\tilde{G}}$. The rest of the vertices in I form a maximal independent set in subgraph, generated by the set

$$L_{\tilde{G}} \cup \{\tilde{u}_i \mid i \leq t\} \cup \{\tilde{v}_i \mid i \leq t\} \cup \{u_{i,j} \mid i \leq s-1, j \leq t\} \cup \{v_{i,j} \mid i \leq s-1, j \leq t+1\}.$$

The number of the latter is 2^{st} , so we have

$$|\mathcal{I}_2| = 2^{st}. \quad (5)$$

3. Let $\mathcal{I}_3 = \{I \in \mathcal{I}_m(G') \mid I \cap L_G = \emptyset, I \cap R_G \neq \emptyset\}$. As in the previous case we have $|\mathcal{I}_3| = 2^{st}$. With (4) and (5) it gives us

$$|\mathcal{I}_1| + |\mathcal{I}_2| + |\mathcal{I}_3| = \iota_m(G) \cdot 2^{st}. \quad (6)$$

4. We now turn to counting those m.i.s. of G' that contain no vertices of V_G . We use the notation $\mathcal{I}_{\overline{G}} = \{I \in \mathcal{I}_m(G') \mid I \cap V_G = \emptyset\}$.

Let $\hat{G}$ be a subgraph of G generated by vertices $u_{i,j}$ and $v_{i,j}$, $1 \leq i \leq s-1$, $1 \leq j \leq t+1$. For what follows it is useful to calculate $\iota_m(\hat{G})$. The number $\hat{\iota}_0$ of m.i.s. of $\hat{G}$ which do not contain any $v_{i,t+1}$ equals to $2^{(s-1)t}$ (that is the number of m.i.s. in a matching with $(s-1)t$ edges). Next consider an arbitrary k , $1 \leq k \leq s-1$. Let us count the number $\hat{\iota}_k$ of those maximal independent sets $\hat{I}$ in $\hat{G}$, that contain $v_{k,t+1}$ but do not contain any of $v_{i,t+1}$ for $i > k$. For such $\hat{I}$ we have $\hat{I} \not\ni u_{i,j}$ and $\hat{I} \ni v_{i,j}$ for all $i < k$ and for all j . Moreover, for such $\hat{I}$ we have $\hat{I} \ni u_{i,t+1}$ for $i > k$, and the rest of vertices in $\hat{I}$ form a maximal independent set in a matching $\{u_{i,j}v_{i,j} \mid k < i \leq s-1, 1 \leq j \leq t\}$. From what was mentioned it follows that $\hat{\iota}_k = 2^{(s-1-k)t}$. Finally we have

$$\iota_m(\hat{G}) = \sum_{k=0}^{s-1} \hat{\iota}_k = \sum_{k=0}^{s-1} 2^{(s-1-k)t} = \frac{2^{st} - 1}{2^t - 1}. \quad (7)$$

- (a) Let $\mathcal{I}_4 = \{I \in \mathcal{I}_{\overline{G}} \mid I \cap L_{\tilde{G}} \neq \emptyset, I \cap R_{\tilde{G}} \neq \emptyset\}$. For $I \in \mathcal{I}_4$ the subset $I \cap V_{\tilde{G}}$ is a m.i.s. in $\tilde{G}$, and the set $I \cap (V_G \setminus V_{\tilde{G}})$ is a m.i.s. in $\hat{G}$. So we get

$$|\mathcal{I}_4| = (\iota_m(\tilde{G}) - 2) \cdot \iota_m(\hat{G}). \quad (8)$$

- (b) Let $\mathcal{I}_5 = \{I \in \mathcal{I}_{\overline{G}} \mid I \cap L_{\tilde{G}} \neq \emptyset, I \cap R_{\tilde{G}} = \emptyset\}$. Every $I \in \mathcal{I}_5$ contains all vertices of $L_{\tilde{G}}$ and every $\tilde{u}_i$. Moreover, such I would not contain w and any $\tilde{v}_i$. Note that $I \cap V_{\tilde{G}}$ is a m.i.s. in $I \cap V_{\tilde{G}}$ and should contain at least one of the vertices $v_{i,t+1}$. It implies

$$\mathcal{I}_5 = \iota_m(\hat{G}) - 2^{(s-1)t}. \quad (9)$$

- (c) Let $\mathcal{I}_6 = \{I \in \mathcal{I}_{\overline{G}} \mid I \cap L_{\tilde{G}} = \emptyset, I \cap R_{\tilde{G}} \neq \emptyset\}$. Similar to the previous case we get that for every $I \in \mathcal{I}_6$ the set $I \cap V_{\tilde{G}}$ is a m.i.s. in $I \cap V_{\tilde{G}}$ and should contain at least one of the vertices $u_{i,t+1}$. So we have

$$\mathcal{I}_6 = \iota_m(\hat{G}) - 1. \quad (10)$$

- (d) It now suffices to find the size of $\mathcal{I}_7 = \{I \in \mathcal{I}_{\overline{G}} \mid I \cap V_{\tilde{G}} = \emptyset\}$. For every $I \in \mathcal{I}_7$ we have $w \in I$. The set $\tilde{I} = I \cap (\{\tilde{u}_i \mid i \leq t\} \cup \{\tilde{v}_i \mid i \leq t\})$ should contain at least one of $\tilde{v}_i$ and should be a m.i.s. in the corresponding subgraph. The number of such $\tilde{I}$ equals to $(2^t - 1)$. The set $\hat{I} = I \setminus (\{w\} \cup \tilde{I})$ should be a m.i.s. in $\hat{G}$ and should contain at least one of $v_{i,t+1}$. The number of choices for such $\hat{I}$ is $(\iota_m(\hat{G}) - 2^{(s-1)t})$. At last we have

$$|\mathcal{I}_7| = (2^t - 1)(\iota_m(\hat{G}) - 2^{(s-1)t}). \quad (11)$$

By (6), (8), (9), (10), (11) and (7), after some calculations we get

$$\iota_m(G') = \sum_{k=1}^7 |\mathcal{I}_k| = 2^{st} \cdot \iota_m(G) + \frac{2^{st} - 1}{2^t - 1} \cdot \iota_m(\tilde{G}) - 2.$$

It suffices to apply lemma 4 to G' . □

Let $\bar{n}$ denote the binary representation of n . Let $w^{(k)}$ denote a binary word which consists of word w repeated k times.

Lemma 7. *Let $n, p, q \in \mathbb{N}$, $n \geq 2$. Let n' be a natural number with binary representation $\bar{n}w^{(q)}$, where w is a binary word of length p . Let G be bipartite without isolated vertices, having $\iota_m(G) = n$. Then there is a bipartite graph without isolated vertices having n' maximal independent sets and no more than $\nu(G) + 2pq + 20(p + \sqrt{pq})$ vertices.*

Proof. If w contains only zeros, then the desired graph can be obtained by adding a matching on $2pq$ vertices to G . For the rest of the proof we assume w being a not-all-zero word. Firstly we consider the case $q = 1$. If $w = 0 \dots 01$, then the desired graph is obtained from G by adding a matching on $2(p - 1)$ vertices and applying lemma 3. Otherwise, let $\tilde{n}$ be the number with binary representation w ($\tilde{n} > 1$). Then by a remark to theorem 2, there exists bipartite $\tilde{G}$ without isolated vertices having $\iota_m(\tilde{G}) = \tilde{n}$ and $\nu(\tilde{G}) < 3p$. By applying lemma 6 to G and $\tilde{G}$ with $t = p$ and $s = 1$, we obtain the graph needed.

For the rest of the proof we assume that $q \geq 2$ and w is not an all-zero word. Put $k = \max\{\lceil \sqrt{q/p} \rceil, 2\}$, and let r be the residue of q modulo k . The remark to the theorem 2 implies that there exists $\tilde{G}$ such that $\nu(\tilde{G}) < 3pk$ and the binary representation of $\iota_m(\tilde{G})$ is $w^{(k)}$ with leading zeros trimmed. The application of lemma 6 to G and $\tilde{G}$ with $t = pk$ and $s = \lfloor q/k \rfloor$ gives us a graph G'' with the binary representation of $\iota_m(G'')$ being $\bar{n}w^{(q-r)}$ and

$$\nu(G'') \leq \nu(G) + 3pk + 2(q/k)(pk + 1) + 3 = \nu(G) + 2pq + 3pk + 2q/k + 3. \quad (12)$$

This, together with the inequalities $2q/k \leq 2\sqrt{pq}$ and $k \leq 2 + \sqrt{q/p}$ implies

$$\nu(G'') \leq \nu(G) + 2pq + 6p + 5\sqrt{pq} + 3.$$

If $r = 0$, then G'' is the desired graph. If $r > 0$, then using the remark to theorem 2, consider a graph $\tilde{G}_r$ having $\nu(\tilde{G}_r) \leq 3r$, and the binary representation of $\iota_m(\tilde{G}_r)$ being equal to $w^{(r)}$ with leading zeros trimmed. Then, by lemma 6 (applied with G'' and $\tilde{G}_r$ as graph G and $\tilde{G}$ respectively, $s = 1$ and $t = pr$), there exists G' having $\overline{\iota_m(G')} = \overline{\iota_m(G'')}w^{(pr)} = \bar{n}'$ and

$$\nu(G') \leq \nu(G'') + 3r + 2pr + 5 \leq \nu(G) + 2pq + 6p + 5\sqrt{pq} + 3r + 2pr + 8.$$

Using the inequality $r < k \leq 2 + \sqrt{q/p}$, we get

$$\nu(G') \leq \nu(G) + 2pq + 10p + 9\sqrt{pq} + 8 < \nu(G) + 2pq + 20(p + \sqrt{pq}).$$

□

Theorem 3. *Let n be a natural number with its binary representation of the form $w_1^{(q_1)} \dots w_k^{(q_k)}$. Let p_i be the length of w_i . If $\sum_{i=1}^k p_i = o(\log n)$ then the following asymptotic holds for arbitrary q_i :*

$$L_{\iota_m, \nu}^{\mathcal{B}}(n) \sim 2 \log_2 n. \quad (13)$$

Proof. The lower bound was already stated in theorem 2, so we proceed to the upper. Lemma 7 implies that there is a graph G with $\iota_m(G) = n$ and

$$\nu(G) \leq 2 \log_2 n + O\left(\sum_{i=1}^k p_i + \sum_{i=1}^k \sqrt{p_i q_i}\right). \quad (14)$$

Then the inequality $\sum_{i=1}^k p_i q_i < 2 \log_2 n$ and Cauchy–Bunyakovsky–Schwarz inequality imply

$$\sum_{i=1}^k \sqrt{p_i q_i} \leq \sqrt{2k \log_2 n} = o(\log n). \quad (15)$$

Finally (14) and (15) imply (13). □

The research was supported by the Russian President Grant MK-3429.2010.1 and RFBR Grant No.10-01-00768a.

References

- [1] Czabarka E., Szekely L., Wagner S. The inverse problem for certain tree parameters // Discrete Applied Mathematics. — 2009. 15 (157). — P. 3314–3319.
- [2] Erdős P., Gallai T. Grafok előirt fokú pontokkal // Mat. Lapok 11 (1960), pp. 264–274.
- [3] Havel V. Poznámka o existenci konečných grafů // Časopis pro pěstování matematiky, Vol. 80 (1955), No. 4, 477–480.
- [4] Linek V. Bipartite graphs can have any number of independent sets // Discrete Mathematics. — 1989. 2 (76). — P. 131–136.
- [5] Wagner S. A class of trees and its Wiener index // Acta Appl. Math. 91 (2) (2006) P. 119–132.
- [6] Wang H., Yu G. All but 49 numbers are Wiener indices of trees // Acta Appl. Math. 92 (1) (2006) P. 15–20.